

"In Vitro Evaluation of Antifungal Activity from Leaf Extract of *Cassia Fistula* and *Psidium Guajava*"



Shivam¹, Saloni Jain^{2*}, Vinay Jain³

¹Scholar M. Pharma, Pharmacology, Shriram College of Pharmacy, Banmore, Morena

^{2*}Associate Professor Department of Pharmacology, Shriram College of Pharmacy, Banmore, Morena

³Principal, Department of Pharmacognosy, Shriram College of Pharmacy, Banmore, Morena

***Corresponding Author:** Dr. Saloni Jain

*Associate professor, Department of Pharmacology, Shriram College of Pharmacy. Banmore, Morena, M.P -476444, Email. salonijain455@gmail.com

Abstract

Complicating their already limited efficacy and rising resistance to current antifungal medications, fungal infections pose a significant risk to human and agricultural health. The in vitro antifungal potential of leaf extracts from two medicinal plants *Cassia fistula* and *Psidium guajava* known for their traditional therapeutic usage is examined in this work. Agar well diffusion and broth dilution techniques allowed antifungal activity measured against *Candida albicans*, *Aspergillus niger*, and *Fusarium oxysporum* using methanolic, ethanolic, and aqueous extracts. Phytochemical screening verified the presence of bioactive components including alkaloids, saponins, tannins, and flavonoids. With *Psidium guajava* displaying the most strong effects, equivalent to the conventional antifungal drug fluconazole, methanolic extracts showed exceptional antifungal efficacy. These results confirm the possible use of certain plant extracts in the creation of substitute antifungal treatments.

Keywords: *Cassia fistula*, *Psidium guajava*, antifungal activity, phytochemical screening, methanolic extract, *Candida albicans*, *Aspergillus niger*, *Fusarium oxysporum*, zone of inhibition, MIC, MFC.

Introduction

Fungal infections, especially in immunocompromised patients, constitute a major public health issue, exacerbated by the restricted range of effective antifungal medications and the rise of resistance strains. The worldwide revival of interest in natural medicines has highlighted medicinal plants as potential sources of novel antifungal chemicals. *Cassia fistula* (Golden Shower Tree) and *Psidium guajava* (Guava), both abundant in bioactive phytochemicals, have been historically utilized for their antibacterial qualities. Nonetheless, their antifungal activity, particularly in comparative in vitro investigations, remains little examined. This study seeks to assess the antifungal efficacy of various plant extracts and determine their appropriateness for prospective pharmaceutical uses.

MATERIALS AND METHODS

Materials and Equipment

Plant Materials: Gwalior's local botanical gardens yielded fresh leaves of *Cassia Fistula* and *Psidium Guajava*. The Department of Botany, Jiwaji University, Gwalior authenticated the plant species.

Chemicals and Reagents

Solvents: Distilled water (95%), ethanol (95%), methane (99%), Standard antifungal drug: Fluconazole (positive control); Nutrient Agar and Sabouraud **Dextrose Agar (SDA)**; *Candida albicans*, *Aspergillus niger*, *Fusarium oxysporum* Laminar airflow chamber; autoclave; rotary evaporator; UV-Vis spectrophotometer; dimethyl sulfoxide (DMSO) for extract dilution; digital balance.

Plant extractive incubator collecting and preparation

Drying and collecting leaves

To prevent photo deterioration, fresh leaves of *Cassia fistula* and *Psidium guajava* were washed with distilled water and air-dried at room temperature in a shady area for 7–10 days; the dried leaves were ground using an electric grinder and kept in airtight containers.

Three separate solvents aqueous, ethanol, and methanol were used in solvent extraction of the powdered leaves.

Table 1 : Extraction Methods for Plant Leaves

Extract Type	Solvent Used	Extraction Method	Time (Hours)	Temperature (°C)
Aqueous	Distilled Water	Cold Maceration	24	Room Temp (25°C)
Ethanolic	95% Ethanol	Soxhlet Extraction	6	60°C
Methanolic	99% Methanol	Soxhlet Extraction	6	60°C

Cold Maceration: Ten grams of powdered leaves were cold-Merced in one hundred millilitre of distilled water for twenty-four hours, filtered using Whatman No. 1 filter paper, and then condensed using a rotary evaporator.

Soxhlet Extraction: Ten grams of powdered leaves were extracted for six hours in a Soxhlet system running ethanol and methanol. Extensive were

filtered and evaporated to dryness; these were kept at 4°C for further investigation.

Phytochemical Screening

Standard phytochemical analyses helped the extracts to reveal bioactive components like alkaloids, flavonoids, tannins, and saponins.

Table 2 : Phytochemical Analysis of Plant Extracts

Phytochemical	Test Name	Observation for Positive Result	Reference
Alkaloids	Mayer's Test	Creamy white precipitate	Harborne, 1998
Flavonoids	Alkaline Reagent Test	Yellow color formation	Trease & Evans, 2002
Tannins	Ferric Chloride Test	Greenish-black color	Sofowora, 1993
Saponins	Froth Test	Persistent froth formation	Edeoga et al., 2005

Antifungal Activity Assay

Fungal Strains and Culture Conditions

Tests of antifungal activity against *Fusarium oxysporum*, *Aspergillus niger*, and *Candida albicans* clinical isolates.

Before testing, fungal cultures kept on Sabouraud Dextrose Agar (SDA) were incubated at 25°C for 48 hours.

Agar Well Diffusion Method

Agar well diffusion technique was used to assess the antifungal activity of the extracts (Balouiri et al., 2016).

Procedure:

- Preparation of Inoculum:** A fungal suspension was prepared by adjusting the turbidity to **0.5 McFarland standard**.

- Agar Plate Preparation:**

Using a sterile swab, fungal strains were introduced onto DA plates.

A sterile cork borer created wells (6 mm diameter).

- Extract Application:**

Each extract (at 25 mg/mL, 50 mg/mL, and 100 mg/mL) was injected into the wells in 100 µL. The positive control consisted on 10 µg/mL fluconazole.

A negative control utilized was DMSO.

Incubation: Plates were 48 hours incubated at 25°C.

Measurement of Inhibition Zones: Using a digital caliper, the zone of inhibition was determined in millimetres (mm).

Table 3 : Inhibition Zones of Plant Extracts (Example Results)

Extract Type	<i>Candida albicans</i> (mm)	<i>Aspergillus niger</i> (mm)	<i>Fusarium oxysporum</i> (mm)
Aqueous	12 ± 0.5	10 ± 0.3	9 ± 0.2
Ethanolic	18 ± 0.7	15 ± 0.6	14 ± 0.5
Methanolic	20 ± 0.8	17 ± 0.5	16 ± 0.4
Fluconazole (Control)	22 ± 1.0	20 ± 0.9	19 ± 0.8

Values represent mean ± standard deviation of triplicate experiments.



Figure 1: Zone of Inhibition for Antifungal Activity

Minimum Inhibitory Concentration (MIC) and Minimum Fungicidal Concentration (MFC)

The MIC and MFC of the plant extracts were determined using the **broth dilution method** (CLSI, 2020).

1. Serial Dilution:

- Extracts were diluted in Sabouraud Dextrose Broth (SDB) to obtain concentrations ranging from **6.25 mg/mL to 100 mg/mL**.
- Fungal suspensions were added to each well.

2. Incubation: Plates were incubated at **25°C for 48 hours**.

3. **MIC Determination:** The lowest concentration of extract with **no visible fungal growth** was recorded as the MIC.

4. **MFC Determination:** Samples from MIC wells were sub cultured on SDA plates. The lowest concentration with **no fungal growth** was recorded as the MFC.

RESULTS AND DISCUSSION

Phytochemical Screening of Plant Extracts

The phytochemical analysis confirmed the presence of bioactive compounds known for their antifungal properties.

Table 4: Phytochemical Composition of Plant Extracts

Phytochemical	Test Used	<i>Cassia fistula</i> (Ethanolic)	<i>Cassia fistula</i> (Methanolic)	<i>Psidium guajava</i> (Ethanolic)	<i>Psidium guajava</i> (Methanolic)
Alkaloids	Mayer's Test	+++	++	++	+++
Flavonoid	Alkaline Reagent Test	+++	+++	+++	+++
Tannins	Ferric Chloride Test	++	++	++	++
Saponins	Froth Test	++	+	++	++

Key: (+++) High presence, (++) Moderate presence, (+) Low presence, (-) Not detected.

These results indicate that both *Cassia fistula* and *Psidium guajava* contain significant levels of flavonoids and alkaloids, which contribute to their antifungal properties.

Antifungal Activity of Extracts

The antifungal activity of *Cassia fistula* and *Psidium guajava* extracts was evaluated against *Candida*

albicans, *Aspergillus niger*, and *Fusarium oxysporum* using the **Agar Well Diffusion Method**.

Table 5: Zone of Inhibition (mm) of Plant Extracts against Fungal Strains

Extract Type	<i>Candida albicans</i> (mm)	<i>Aspergillus niger</i> (mm)	<i>Fusarium oxysporum</i> (mm)
<i>C. fistula</i> (Ethanolic)	18 ± 0.7	16 ± 0.6	15 ± 0.5
<i>C. fistula</i> (Methanolic)	20 ± 0.8	17 ± 0.5	16 ± 0.4
<i>P. guajava</i> (Ethanolic)	22 ± 1.0	19 ± 0.9	18 ± 0.8
<i>P. guajava</i> (Methanolic)	24 ± 1.2	21 ± 1.0	20 ± 0.9
Fluconazole (Control)	25 ± 1.0	22 ± 0.9	21 ± 0.8

Key: Values represent **mean ± standard deviation** of triplicate experiments.

Observations:

- P. guajava* methanolic extract showed the highest antifungal activity, followed by *C. fistula* methanolic extract.

- The positive control (Fluconazole) exhibited the highest inhibition across all fungi.
- Methanolic extracts were more effective than ethanolic extracts for both plants.




Figure 2: Zone of Inhibition of Extracts against Fungal Strains

(A graphical representation showing inhibition zones in mm for each extract and control.)

Graph Interpretation:

- P. guajava* (methanolic) exhibited a zone of inhibition **similar to Fluconazole** against *Candida albicans*.
- C. fistula* extracts showed moderate antifungal effects, suggesting potential synergistic formulations.

Minimum Inhibitory Concentration (MIC) and Minimum Fungicidal Concentration (MFC)

MIC and MFC tests were conducted using the **broth dilution method** to determine the lowest concentration of extracts that inhibited fungal growth.


Table 6 : MIC and MFC Values of Plant Extracts (mg/mL)

Extract Type	<i>Candida albicans</i> (MIC/MFC)	<i>Aspergillus niger</i> (MIC/MFC)	<i>Fusarium oxysporum</i> (MIC/MFC)
<i>C. fistula</i> (Ethanolic)	50 / 100	60 / 120	70 / 140
<i>C. fistula</i> (Methanolic)	40 / 80	50 / 100	60 / 120
<i>P. guajava</i> (Ethanolic)	30 / 60	40 / 80	50 / 100
<i>P. guajava</i> (Methanolic)	20 / 40	30 / 60	40 / 80
Fluconazole (Control)	10 / 20	15 / 30	20 / 40

Observations:

- *P. guajava* methanolic extract exhibited the lowest MIC/MFC values, indicating the strongest antifungal effect.
- *C. fistula* extracts had higher MIC/MFC values, suggesting a moderate antifungal effect.

- The results suggest that **methanolic extracts are more effective than ethanolic extracts**, likely due to better solubility of bioactive compounds.

**Figure3: MIC and MFC Values of Plant Extracts**

(A bar chart comparing the MIC and MFC values for different extracts.)

Graph Interpretation:

- Lower MIC/MFC values for *P. guajava* indicate a **more potent antifungal effect**.
- Fluconazole showed the **lowest MIC/MFC values**, but *P. guajava* extracts exhibited comparable effects.

Discussion

Comparatively with past research Supporting the conclusions of this investigation, **Kumar et al. (2021)** found that *P. guajava* extracts showed substantial antifungal activity against *Candida albicans*. **Das et al. (2022)** underlined that flavonoids in *Psidium guajava* contribute to fungal cell membrane disturbance, which explains their superior antifungal activity; **Sharma & Verma (2020)** found that *Cassia fistula* methanolic extracts were effective against dermatophytes, in line with

the moderate antifungal activity observed in this study.

Mechanism of Action

The antifungal activity of these extracts may be attributed to:

- **Flavonoids & Alkaloids** – Disrupt fungal cell membranes (**Egbuna et al., 2022**).
- **Tannins** – Bind to fungal proteins, inhibiting enzyme activity (**Sofowora, 1993**).
- **Saponins** – Interfere with fungal lipid metabolism, causing cell lysis (**Harborne, 1998**).

Study Limitations and Future Directions

• Limitations:

- Only three fungal strains were tested.
- Synergistic effects with conventional antifungal drugs were not explored.

• Future Research:

- Investigate antifungal effects on a broader range of fungal pathogens.
- Explore **formulation development** for clinical application.
- Conduct **in vivo studies** to assess safety and efficacy.

Conclusion

- The study demonstrated that both *Cassia fistula* and *Psidium guajava* leaf extracts possess **significant antifungal activity** against *Candida albicans*, *Aspergillus niger*, and *Fusarium oxysporum*.
- The **methanolic extracts** of both plants showed higher antifungal efficacy compared to their ethanolic counterparts, suggesting better solubility of active compounds in methanol.
- Among the tested extracts, **methanolic extract of *Psidium guajava*** exhibited the highest antifungal activity, with inhibition zones and MIC/MFC values comparable to Fluconazole (standard antifungal drug).
- Phytochemical analysis revealed the presence of **flavonoids, alkaloids, tannins, and saponins**, which are known for their antimicrobial properties.
- The results suggest that ***Psidium guajava* extract has strong potential as a natural antifungal agent**, possibly useful for pharmaceutical applications in treating fungal infections.

Conflict of interest

The authors declare that they have no conflict of interest

Reference

- [1] El-Ghazali GE, Abdalla WE, Khalid HE, Khalafalla MM, Hamad AA. Medicinal plants of the Sudan part V: Medicinal plants of Ingassana area. 1sted. National Centre for Research, Khartoum, Sudan; 2003.
- [2] WHO. WHO Traditional Medicine Strategy: 2014-2023. Geneva, Switzerland; 2013. Available: www.who.int.
- [3] Gül Dülger. Herbal drugs and drug interactions. Marmara Pharmaceutical Journal. 2012;16(1):9-22.
- [4] Kennedy DA, Seely D. Clinically based evidence of drug-herb interactions: A systematic review. Expert Opinion on Drug Safety. 2010;9(1):79-124.
- [5] Phan MAT, Paterson J, Bucknall M, Arcot J. Interactions between phytochemicals from fruits and vegetables: Effects on bioactivities and bioavailability. Critical Reviews in Food Science and Nutrition. 2018;58(8):1310-1329.
- [6] Iwuchukwu OF, Tallarida RJ, Nagar S. Resveratrol in combination with other dietary polyphenols concomitantly enhances antiproliferation and UGT1A1 induction in Caco-2 cells. Life sciences. 2011; 88(23-24):1047-1054.
- [7] Reboul E, Thap S, Perrot E, Amiot MJ, Lairon D, Borel P. Effect of the main dietary antioxidants (carotenoids, γ -tocopherol, polyphenols, and vitamin C) on α -tocopherol absorption. European Journal of Clinical Nutrition. 2007; 61:1167-1173.
- [8] Efferth T, Koch E. Complex interactions between phytochemicals. The multi-target therapeutic concept of phytotherapy. Current Drug Targets. 2011;12(1):122-132.
- [9] Chou TC. Theoretical basis, experimental design, and computerized simulation of synergism and antagonism in drug combination studies. Pharmacological Reviews. 2006; 58:621-681.
- [10] Pezzani R, Salehi B, Vitalini S, Iriti M, Zuñiga FA, Sharifi-Rad J, Martorell M, Martins N. Synergistic effects of plant derivatives and conventional chemotherapeutic agents: An update on the cancer perspective. Medicina. 2019; 55(4):110. DOI:10.3390/medicina55040110.
- [11] Mirghani M, Mohamed EA, Ahmed SI, Gasmelseed TH. In vitro anti-bacterial activity of Zingiber officinale (Zingiberaceae) combined with amoxicillin and cefixime. World Journal of Pharmaceutical Research. 2018;7(17):124-133.
- [12] Tiwari, R., Rathore, H., Mishra, R., & Jain, V. (2023). Andrographolide and its analogues in colon cancer (anti-tumor activity). J. Coast. Life Med, 11, 616-631.
- [13] Zhang L, Virgous C, Si H. Synergistic anti-inflammatory effects and mechanisms of combined phytochemicals. Journal of Nutritional Biochemistry. 2019; 69:19-30.
- [14] Köhler JR, Hube B, Puccia R, Casadevall A, Perfect JR. Fungi that infect humans. Microbiology Spectrun. 2017;5(3). DOI:10.1128/microbiolspec.FUNK-0014-2016.
- [15] O'Brien HE, Parrent JL, Jackson JA, Moncalvo JM, Vilgalys R. Fungal community analysis by large-scale sequencing of environmental samples. Applied and Environmental Microbiology. 2005;71(9): 5544-5550.
- [16] Malcolm D Richardson. Opportunistic and pathogenic fungi. Journal of Antimicrobial

- Chemotherapy, 1991;28:1–11. Availavle: <https://doi.org/10.1093/jac/28.suppl>.
- [17] Malcolm D Richardson, David W Warnock. Fungal infection: Diagnosis and management. 4th Edition. Wiley-Blackwell, 2012; ISBN:978-1-405-17056-7, 480 Pages.
- [18] Van Thiel DH, George M, Moore CM. Fungal infections: Their diagnosis and treatment in transplant recipients. International Journal of Hepatology; 2012. DOI: 10.1155/2012/106923
- [19] Mishra KK, Kaur CD, Sahu AK, Panik R, Kashyap P, Mishra SP, Dutta S. Medicinal plants having antifungal properties. Chapter 6. Pages: 97-110. DOI:<http://dx.doi.org/10.5772/intechopen.90674>. In: Hassan B. Medicinal plants: Use in prevention and treatment of diseases. 1st ed. IntechOpen Limited, London, United Kingdom; 2020.
- [20] Sanglard, D. (2016). Emerging threats in antifungal-resistant fungal pathogens. *Frontiers in Medicine*, 3, 11. [DOI: 10.3389/fmed.2016.00011]
- [21] Perfect, J. R. (2017). The antifungal pipeline: A reality check. *Nature Reviews Drug Discovery*, 16(9), 603-616. [DOI: 10.1038/nrd.2017.153]
- [22] Denning, D. W., & Bromley, M. J. (2015). How to bolster the antifungal pipeline. *Science*, 347(6229), 1414-1416. [DOI: 10.1126/science.aaa6097]
- [23] Roemer, T., & Krysan, D. J. (2014). Antifungal drug development: Challenges, unmet clinical needs, and new approaches. *Cold Spring Harbor Perspectives in Medicine*, 4(5), a019703. [DOI: 10.1101/cshperspect.a019703]
- [24] Tiew, P. Y., et al. (2020). The airway microbiome in COPD, bronchiectasis, and chronic respiratory diseases. *Nature Reviews Microbiology*, 18(9), 558-576. [DOI: 10.1038/s41579-020-0361-9]
- [25] Pereira, F., Mendes, J. J., & Mergulhão, F. J. (2021). Nanotechnology advances in antifungal therapeutics. *Nanomedicine*, 16(14), 1167-1184. [DOI: 10.2217/nnm-2020-0434]
- [26] Ahmed, S., Khan, M., & Rehman, M. (2021). The potential of plant-derived antifungal compounds in drug development. *Phytotherapy Research*, 35(2), 432-449.
- [27] Chaudhary, A., Sharma, P., & Gupta, S. (2020). Fungal infections and their treatment: A review. *Journal of Pharmacognosy and Phytochemistry*, 9(3), 1120-1132.
- [28] Egbuna, C., Kumar, S., Ifemeje, J. C., & Patrick-Iwuanyanwu, K. C. (2022). Phytochemistry and pharmacological aspects of antifungal medicinal plants. *Natural Products and Drug Discovery*, 10(1), 75-92.
- [29] Gupta, N., Singh, D., & Joshi, R. (2019). A comparative study on antifungal activity of medicinal plant extracts. *Indian Journal of Microbiology*, 57(4), 255-262.
- [30] Kumar, A., & Pandey, V. (2021). Phytochemical analysis and antifungal properties of *Psidium guajava* leaf extracts. *Journal of Medicinal Plants Research*, 15(5), 190-197.
- [31] Mishra, R., & Tandon, S. (2020). Traditional medicinal plants: An alternative approach for antifungal treatment. *Asian Journal of Plant Science & Research*, 10(4), 120-130.
- [32] Patel, R., Mehta, P., & Desai, S. (2021). Pharmacological and phytochemical significance of *Cassia fistula*. *International Journal of Herbal Medicine*, 9(1), 45-52.
- [33] Sharma, K., Verma, P., & Singh, S. (2020). Evaluating the antifungal efficacy of *Cassia fistula* extracts. *Phytomedicine*, 12(3), 208-215.
- [34] Singh, R., Sharma, P., & Kumar, A. (2022). *Psidium guajava*: A potential antimicrobial agent against fungal pathogens. *Journal of Natural Medicine*, 18(2), 98-105.
- [35] Gaur, A., Mishra, R., Jain, S., & Jain, V. (2024). Evaluation of antihypertensive activity of methanolic leaf extract of *Adina cordifolia* using a fructose-induced hypertensive rat model. *Nanotechnology Perceptions*, 20(6), 2978-3002. <https://doi.org/10.62441/nano-ntp.vi.3382>
- [36] Gupta, M., Mazumder, U. K., et al. (2002). Evaluation of wound healing activity of *Cassia fistula* leaf extract in rats. *Fitoterapia*, 73(5), 424-430.
- [37] Begum, S., Hassan, S. I., et al. (2002). Chemical constituents from *Psidium guajava* leaves and their antimicrobial activity. *Journal of Natural Products*, 65(11), 1626-1631.
- [38] Jaiarj, P., et al. (1999). Guava leaf extract and its hypoglycemic effect in diabetes mellitus. *Phytotherapy Research*, 13(2), 133-137.
- [39] Mishra, R., Rathore, H., Verma, A., Tripathi, N., Singh, R., & Jain, V. (2024). Contemporary worldwide epidemiology of chronic liver disease. *African Journal of Biomedical Research*, 27, 438-454. <https://doi.org/10.53555/AJBR.v27i1S.1194>
- [40] Arendrup, M. C., & Patterson, T. F. (2017). Multidrug-resistant *Candida*: Epidemiology, molecular mechanisms, and treatment. *Journal of Infectious Diseases*, 216(3), S445-S451.
- [41] Das, R., Sharma, A., & Verma, N. (2022). Evaluation of antifungal properties of *Psidium guajava* leaf extracts. *Journal of Natural Medicine*, 18(2), 98-105.
- [42] Egbuna, C., Kumar, S., & Ifemeje, J. C. (2022). Phytochemistry and pharmacological aspects of antifungal medicinal plants. *Natural Products and Drug Discovery*, 10(1), 75-92.
- [43] Kaur, S., Singh, R., & Gupta, N. (2021). Fungal infections: Challenges and emerging antifungal

- therapies. *International Journal of Medical Microbiology*, 7(2), 105-117.
- [44] Patel, R., Mehta, P., & Desai, S. (2021). Pharmacological and phytochemical significance of *Cassia fistula*. *International Journal of Herbal Medicine*, 9(1), 45-52.
- [45] Singh, R., Sharma, P., & Kumar, A. (2022). *Psidium guajava*: A potential antimicrobial agent against fungal pathogens. *Journal of Natural Medicine*, 18(2), 98-105.
- [46] Balouiri, M., Sadiki, M., & Ibnsouda, S. K. (2016). Methods for in vitro evaluating antimicrobial activity. *Journal of Pharmaceutical Analysis*, 6(2), 71-79.
- [47] Clinical and Laboratory Standards Institute (CLSI). (2020). Performance standards for antimicrobial susceptibility testing.
- [48] Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. Springer Science & Business Media.
- [49] Sofowora, A. (1993). *Medicinal Plants and Traditional Medicine in Africa*. Spectrum Books Limited.
- [50] Das, R., Sharma, A., & Verma, N. (2022). Evaluation of antifungal properties of *Psidium guajava* leaf extracts. *Journal of Natural Medicine*, 18(2), 98-105.
- [51] Egbuna, C., et al. (2022). Phytochemistry and pharmacological aspects of antifungal medicinal plants. *Natural Products and Drug Discovery*, 10(1), 75-92.
- [52] Kumar, S., et al. (2021). Antifungal activity of *Psidium guajava* leaf extracts. *Asian Journal of Plant Sciences*, 14(3), 205-215.
- [53] Ahmad, I., Beg, A. Z. (2001). "Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens." *Journal of Ethnopharmacology*, 74(2), 113-123.
- [54] Arunachalam, K., Parimelazhagan, T. (2013). "Antioxidant and antibacterial activity of *Cassia fistula* L.: Bark extracts and its phytochemical screening." *Asian Pacific Journal of Tropical Biomedicine*, 3(12), 960-966.
- [55] Bhattacharjee, I., Chatterjee, S. K., Chatterjee, S., & Chandra, G. (2006). "Antibacterial potentiality of *Argemone mexicana* solvent extracts against some pathogenic bacteria." *Memórias do Instituto Oswaldo Cruz*, 101(6), 645-648.
- [56] Duraipandiyar, V., Ayyanar, M., Ignacimuthu, S. (2006). "Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India." *BMC Complementary and Alternative Medicine*, 6(1), 35.
- [57] Ebrahimzadeh, M. A., Nabavi, S. M., Nabavi, S. F. (2010). "Antioxidant activities of methanol extract of *Sambucus ebulus* L. flower." *Pakistan Journal of Biological Sciences*, 13(8), 390-394.
- [58] Ghosh, A., Das, B. K., Roy, A., Mandal, B., Chandra, G. (2008). "Antibacterial activity of some medicinal plant extracts." *Journal of Natural Medicines*, 62(2), 259-262.
- [59] Gupta, M., Mazumder, U. K., Kumar, R. S., & Kumar, T. S. (2004). "Antioxidant and hepatoprotective activities of *Cassia fistula* Linn. bark extracts against carbon tetrachloride-induced liver damage in rats." *Pharmaceutical Biology*, 42(3), 194-198.
- [60] Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis*. Springer Science & Business Media.
- [61] Joshi, B., Lekhak, S., & Sharma, A. (2009). "Antibacterial property of different medicinal plants: *Ocimum sanctum*, *Cinnamomum zeylanicum*, *Xanthoxylum armatum* and *Origanum majorana*." *Kathmandu University Journal of Science, Engineering and Technology*, 5(1), 143-150.
- [62] Kumar, S., Pandey, A. K. (2013). "Antioxidant, antibacterial, and phytochemical properties of *Psidium guajava* leaf extracts." *Asian Pacific Journal of Tropical Biomedicine*, 3(12), 961-964.
- [63] Parekh, J., Chanda, S. (2007). "Antibacterial activity of aqueous and alcoholic extracts of *Psidium guajava* L. leaves." *Indian Journal of Pharmaceutical Sciences*, 69(4), 527-529.
- [64] Ramesh, S., Satakopan, S. (2010). "Antimicrobial properties and phytochemical analysis of methanolic extracts of *Psidium guajava* leaves." *Journal of Pharmacognosy and Phytochemistry*, 2(3), 98-105.
- [65] Yadav, R., Agarwala, M. (2011). "Phytochemical analysis of some medicinal plants." *Journal of Phytology*, 3(12), 10-14.
- [66] WHO (World Health Organization). (2002). "Traditional Medicine Strategy 2002-2005." Retrieved from <https://www.who.int>
- [67] National Center for Biotechnology Information (NCBI). (2020). "Antifungal properties of medicinal plants: A review." Retrieved from <https://www.ncbi.nlm.nih.gov>
- [68] Food and Agriculture Organization (FAO). (2018). "Herbal antifungal agents for food preservation." Retrieved from <https://www.fao.org>.