American Journal of Psychiatric Rehabilitation

Expert Opinion Article

Holistic Assessment of Individualized Homeopathic Therapies in the Management of Cervical Spondylosis: A Prospective Quasi-Experimental Study at a Tertiary Care Centre

Dr. Padmalaya Mallick[1*], Dr. Shalu Munjal[2], Dr. Vinay Kumar[3]

[1*]Ph.D. (Homoeopathy) (scholar), Tantia University, Sri Ganganagar (Raj.).
[2]M.D.(Hom.), Research Supervisor, Professor, Deptt. of Psychiatry, Sri Ganganagar Homoeopathic Medical College, Hospital & Research Institute, Tantia University.
[3]Ph.D. (Homoeopathy),M.D.(Hom.), Professor, Deptt. of Materia Medica, Sri Ganganagar Homoeopathic Medical College, Hospital & Research Institute, Tantia University.

***Corresponding Author-** Dr. Padmalaya Mallick

- Ph.D (Homoeopathy) (scholar), Tantia University, Sri Ganganagar (Raj.).
- Assist. Prof., Deptt. of Surgery, Orissa Medical college of Homoeopathy & Research, Sambalpur, Odisha.

Abstract
Cervical spondylosis—a prevalent cause of chronic neck pain and functional disability—was studied in 100 adults at a tertiary care centre over 18 months. Participants were treated with individualized homeopathic therapy based on miasmatic analysis and constitution, using rigorous assessment tools: Visual Analogue Scale (VAS), Neck Disability Index (NDI), and Universal Goniometer. Substantial improvements in pain, mobility, and disability profiles were statistically validated. This elaborated report supports the scientific integration of homeopathy for chronic musculoskeletal disorders.

Keywords- *Cervical spondylosis, Neck Disability Index, Visual Analogue Scale, homeopathy.*

Introduction

Cervical spondylosis represents degenerative changes in intervertebral discs and cervical vertebrae that manifest as neck pain, stiffness, radiculopathy, and functional impairment. The condition is linked to modern lifestyle factors—prolonged desk work, poor ergonomics, mental stress, and physical inactivity—making it increasingly common even among younger adults. Epidemiological evidence suggests that up to 95% of individuals exhibit radiographic signs by age 65. Complications may include myelopathy and radiculopathy, impacting daily life and mental health.[1][2]



Age group distribution of cervical spondylosis cases.



Distribution of presenting complaints among cervical spondylosis cases.



Gender distribution of cervical spondylosis cases.

Pathophysiology & Risk Factors:
Cervical spondylosis is characterized by disc dehydration, loss of height, osteophyte formation, and nerve compression (myelopathy/ radiculopathy). Risk factors include age, genetics, occupation, trauma, and inflammatory processes.[2]



Improvement in cervical range of motion with homeopathic management.

A Paradigm for Holistic Care:
Homeopathy treats disease as a disruption of vital force, selecting remedies with similimum based on miasmatic and constitutional analysis. This study aims to scientifically assess the efficacy of individualized homeopathic medicines as a safe,

minimally invasive alternative for cervical spondylosis.

Objectives

- To evaluate the effectiveness of individualized homeopathic medicines in cervical spondylosis using validated quantitative and functional outcome measures.
- To analyze demographic, clinical, and miasmatic patterns among patients.
- To statistically validate pre/post-intervention improvements.

Materials and Methods

Study Design:
A prospective, quasi-experimental, single-centre study with pre- and post-intervention comparison. No control group.

Study Setting:
Tertiary care institution. Duration: 18 months.

Sampling:
Purposive sampling. Sample size calculated for a prevalence rate of 1.9%, CI 95%, permissible error 5%.

Participants:

- Age 21–70 years
- Clinically and radiologically confirmed cervical spondylosis (X-ray)
- Consent for participation and follow-up

Exclusion:

- Traumatic neck injury, whiplash, capsulitis, spinal cord injury, prolapsed disc, pregnancy/lactation, active inflammatory/systemic disease.

Case Documentation:
Comprehensive case-taking included chronology of presenting complaints, past/family history, personal history, detailed physical/systemic examination, and investigations (AP/Lateral X-ray).

Prescription Protocol:
Remedy selection based on the totality of symptoms (mental, physical generals, particulars), repertorial analysis (Kent's, Boenninghausen's, Complete Repertory). Drugs and potency tailored to individual susceptibility, repeated as needed.

Assessment Tools:

- **Visual Analogue Scale (VAS):** Measures pain intensity (0–10; 0–100mm).
- **Neck Disability Index (NDI):** Evaluates disability across 10 daily activities; max score 50.
- **Universal Goniometer:** Assesses cervical range of motion (see normative data in source).

Outcome Categorization:

- *Recovered:* VAS 0–4mm, NDI 0–4, ROM within normal limits.
- *Improved:* VAS 5–74mm, NDI 5–14, moderately improved ROM.
- *Not Improved:* VAS 75–100mm, NDI ≥15, persistent abnormal ROM.

Statistical Analysis:
Mean, SD, CI, paired t-test for pre/post comparisons, significance at $p < 0.05$.

Results

Demographics:

- Highest affected age group: 31–40 years (36%).
- Gender: 53% male, 47% female.
- Occupation: Engineers (22%), teachers (16%), housewives (15%), bank employees (13%).

![Age group distribution of cervical spondylosis cases]
![Gender distribution of cervical spondylosis cases]

Clinical Features:

- Presenting complaints: Neck pain (74%), shoulder pain (37%), headache (26%), stiffness (22%), numbness (20%).
- Miasmatic profiles: Psoro-sycotic (28%), Psora (23%), Sycosis (16%).

![Distribution of presenting complaints among cervical spondylosis cases]

Remedies Prescribed:
Most commonly used: Rhus Toxicodendron (21%), Arnica (15%), Kalmia (13%), Natrum Mur/Calcarea Carb/Sulphur/Paris Quad (10% each).
![Distribution of major homeopathic remedies prescribed in the study]



Distribution of major homeopathic remedies prescribed in the study.

Treatment Outcomes:
Recovered (56%), Improved (33%), Not improved (11%).

![[Clinical outcomes of homeopathic treatment in cervical spondylosis]



Clinical outcomes of homeopathic treatment in cervical spondylosis.

Pain, Disability & ROM Improvements:

- VAS score reduced from mean 7.8±0.57 to 3.2±1.14 ($p = 0.005$).
- NDI decreased from mean 52.8±10.1 to 32.2±10.2.
- Cervical ROM (flexion, rotation) improved substantially:

- Flexion: 45 → 58.3 degrees
- Right rotation: 53.6 → 73 degrees
- Left rotation: 58.6 → 75 degrees

![Reduction in mean pain (VAS) scores after homeopathic treatment]



Reduction in mean pain (VAS) scores after homeopathic treatment.
![Change in Neck Disability Index (NDI) scores before and after homeopathic intervention]



Change in Neck Disability Index (NDI) scores before and after homeopathic intervention.
![Improvement in cervical range of motion with homeopathic management]

Statistical Analysis:
All principal outcome measures showed statistically significant improvements ($p < 0.05$), supporting rejection of the null hypothesis and validating the alternate hypothesis.

Discussion
This study firmly establishes the value of homeopathic therapy for cervical spondylosis. The high rate of positive response (89%) is noteworthy, especially since cases were selected irrespective of chronicity and occupation. The improvement in VAS and NDI aligns with results from previous controlled studies (Nayak et al., Gupta et al.), suggesting homeopathy may effectively target pain, disability, and constitutional factors, avoiding conventional side effects. Miasmatic analysis guided remedy selection, with chronic constitutional remedies predominating, affirming the classical homeopathic approach.
The range of motion data highlights not only symptomatic relief but also functional improvement—crucial for long-term rehabilitation. The inclusion of objective metrics (goniometry, validated scales) strengthens the clinical evidence for homeopathy's effectiveness. Furthermore, treating patients holistically improves their mental health and well-being, often undermined by chronic pain syndromes.
Self-reported outcomes are reinforced by objective data, and advice on posture, ergonomics, and physical exercise supplemented the individualized homeopathic intervention—an integrative strategy for chronic musculoskeletal care.

Limitations
- Lack of a randomized control arm and placebo comparison
- Medium-term follow-up; relapse/recurrence rates not studied
- Single-centre population; results may not generalize universally
- Reliance on self-reported disability for functional assessment

Conclusion
Individualized homeopathic therapy produced statistically and clinically significant improvements in pain, disability, and cervical motion for patients with cervical spondylosis at a tertiary centre. These findings strongly support homeopathy's role as an adjunct in integrated spine care, particularly for populations susceptible to lifestyle-driven degenerative diseases.

Comprehensive References *(sampled; full list available)*
1. Vas T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388:1545-1602.
2. K. Radhakrishna, et al. Epidemiology of cervical radiculopathy: A population-based study. *Brain*. 1994.
3. Shedid D, Benzel EC. Cervical spondylosis: anatomy, pathophysiology, biochemics. *Neurosurgery*. 2007;60(1):S7-13.
4. Andersson GBJ. The epidemiology of spinal disorders. In: Frymoyer JW, ed. *The Adult Spine: Principles and Practice*. Raven Press; 1997:93-141.
5. Naaz Ayesha. Historical overview of cervical spondylosis. *IJRSR*. 2015;6(11):7155-7157.
6. Kudo DT, Tadi P. Cervical Spondylosis. StatPearls [Internet]. 2021.

... (Full reference list up to 70 available, including homeopathy clinical studies, epidemiology, assessment scales, and materia medica sources: Nayak et al., Dsouza et al., Gupta et al., Choudhary et al.)